arXiv:1409.3317v1 [math.NT] 11 Sep 2014

Non-existence of points rational over number fields on Shimura curves

Keisuke Arai

Abstract

Jordan, Rotger and de Vera-Piquero proved that Shimura curves have no points rational over imaginary quadratic fields under a certain assumption. In this article, we expand their results to the case of number fields of higher degree. We also give counterexamples to the Hasse principle on Shimura curves.

1 Introduction

Let B be an indefinite quaternion division algebra over $\mathbb{Q}$, and $d(B)$ its discriminant. Fix a maximal order $\mathcal{O}$ of B. A *QM-abelian surface by* $\mathcal{O}$ over a field F is a pair (A, i) where A is a 2-dimensional abelian variety over F, and $i : \mathcal{O} \hookrightarrow \mathrm{End}_F(A)$ is an injective ring homomorphism satisfying $i(1) = id$ (cf. [2, p.591]). Here, $\mathrm{End}_F(A)$ is the ring of endomorphisms of A defined over F. We assume that A has a left $\mathcal{O}$-action. Let M^B be the Shimura curve over $\mathbb{Q}$ associated to B, which parameterizes isomorphism classes of QM-abelian surfaces by $\mathcal{O}$ (cf. [3, p.93]). We know that M^B is a proper smooth curve over $\mathbb{Q}$. For an imaginary quadratic field k, we have $M^B(k) = \emptyset$ under a certain assumption ([3, Theorem 6.3], [5, Theorem 1.1]). We expand this result to the case of number fields of higher degree in this article. The method of the proof is based on the strategy in [3], and the key is to control the field of definition of the QM-abelian surface corresponding to a rational point on M^B. We also give counterexamples to the Hasse principle on M^B over number fields. We will discuss the relevance to the Manin obstruction in a forthcoming article.

For a prime number q, let $\mathcal{B}(q)$ be the set of isomorphism classes of indefinite quaternion division algebras B over $\mathbb{Q}$ such that

$$\begin{cases} B \otimes_{\mathbb{Q}} \mathbb{Q}(\sqrt{-q}) \ncong \mathrm{M}_2(\mathbb{Q}(\sqrt{-q})) & \text{if } q \neq 2, \\ B \otimes_{\mathbb{Q}} \mathbb{Q}(\sqrt{-1}) \ncong \mathrm{M}_2(\mathbb{Q}(\sqrt{-1})) \text{ and } B \otimes_{\mathbb{Q}} \mathbb{Q}(\sqrt{-2}) \ncong \mathrm{M}_2(\mathbb{Q}(\sqrt{-2})) & \text{if } q = 2. \end{cases}$$

For positive integers N and e, let

$$\mathcal{C}(N, e) := \left\{ \alpha^e + \overline{\alpha}^e \in \mathbb{Z} \;\middle|\; \alpha \in \mathbb{C} \text{ is a root of } T^2 + sT + N \text{ for some } s \in \mathbb{Z},\ s^2 \leq 4N \right\},$$

$$\mathcal{D}(N, e) := \left\{ a, a \pm N^{\frac{e}{2}}, a \pm 2N^{\frac{e}{2}}, a^2 - 3N^e \in \mathbb{R} \;\middle|\; a \in \mathcal{C}(N, e) \right\}.$$

Here, $\overline{\alpha}$ is the complex conjugate of α. If e is even, then $\mathcal{D}(N,e) \subseteq \mathbb{Z}$. For a subset $\mathcal{D} \subseteq \mathbb{Z}$, let

$$\mathcal{P}(\mathcal{D}) := \{ \text{ prime divisors of some of the integers in } \mathcal{D} \setminus \{0\} \}.$$

For a number field k and a prime $\mathfrak{q}$ of k of residue characteristic q, let

- $\kappa(\mathfrak{q})$: the residue field of $\mathfrak{q}$,
- $\mathrm{N}_{\mathfrak{q}}$: the cardinality of $\kappa(\mathfrak{q})$,
- $e_{\mathfrak{q}}$: the ramification index of $\mathfrak{q}$ in $k/\mathbb{Q}$,
- $f_{\mathfrak{q}}$: the degree of the extension $\kappa(\mathfrak{q})/\mathbb{F}_q$,
- $\mathcal{S}(k,\mathfrak{q})$: the set of isomorphism classes of indefinite quaternion division algebras B over $\mathbb{Q}$ such that any prime divisor of $d(B)$ belongs to

$$\begin{cases} \mathcal{P}(\mathcal{D}(\mathrm{N}_{\mathfrak{q}}, e_{\mathfrak{q}})) \cup \{q\} & \text{if } B \otimes_{\mathbb{Q}} k \cong \mathrm{M}_2(k) \text{ and } e_{\mathfrak{q}} \text{ is even,} \\ \mathcal{P}(\mathcal{D}(\mathrm{N}_{\mathfrak{q}}, 2e_{\mathfrak{q}})) \cup \{q\} & \text{if } B \otimes_{\mathbb{Q}} k \ncong \mathrm{M}_2(k). \end{cases}$$

Note that $\mathcal{S}(k,\mathfrak{q})$ is a finite set. The main result of this article is:

Theorem 1.1. *Let k be a number field of even degree, and q a prime number such that*

- *there is a unique prime $\mathfrak{q}$ of k above q,*
- *$f_{\mathfrak{q}}$ is odd (and so $e_{\mathfrak{q}}$ is even), and*
- *$B \in \mathcal{B}(q) \setminus \mathcal{S}(k,\mathfrak{q})$.*

Then $M^B(k) = \emptyset$.

Remark 1.2. (1) By [7, Theorem 0], we have $M^B(\mathbb{R}) = \emptyset$.

(2) If k is of odd degree, then k has a real place, and so $M^B(k) = \emptyset$.

2 Canonical isogeny characters

In this section, we review canonical isogeny characters associated to QM-abelian surfaces, which were introduced in [3, §4]. Let K be a number field, $\overline{K}$ an algebraic closure of K, $\mathrm{G}_K = \mathrm{Gal}(\overline{K}/K)$ the absolute Galois group of K, $\mathcal{O}_K$ the ring of integers of K, (A,i) a QM-abelian surface by $\mathcal{O}$ over K, and p a prime divisor of $d(B)$. Then the p-torsion subgroup $A[p](\overline{K})$ of A has exactly one non-zero proper left $\mathcal{O}$-submodule, which we shall denote by C_p. Then C_p has order p^2, and is stable under the action of G_K. Let $\mathfrak{P}_{\mathcal{O}} \subseteq \mathcal{O}$ be the unique left ideal of reduced norm $p\mathbb{Z}$.

In fact, $\mathfrak{P}_{\mathcal{O}}$ is a two-sided ideal of $\mathcal{O}$. Then C_p is free of rank 1 over $\mathcal{O}/\mathfrak{P}_{\mathcal{O}}$. Fix an isomorphism $\mathcal{O}/\mathfrak{P}_{\mathcal{O}} \cong \mathbb{F}_{p^2}$. The action of G_K on C_p yields a character

$$\varrho_p : \mathrm{G}_K \longrightarrow \mathrm{Aut}_{\mathcal{O}}(C_p) \cong \mathbb{F}_{p^2}^{\times}.$$

Here, $\mathrm{Aut}_{\mathcal{O}}(C_p)$ is the group of $\mathcal{O}$-linear automorphisms of C_p. The character ϱ_p depends on the choice of the isomorphism $\mathcal{O}/\mathfrak{P}_{\mathcal{O}} \cong \mathbb{F}_{p^2}$, but the pair $\{\varrho_p, (\varrho_p)^p\}$ is independent of this choice. Either of the characters $\varrho_p, (\varrho_p)^p$ is called a *canonical isogeny character* at p. We have an induced character

$$\varrho_p^{\mathrm{ab}} : \mathrm{G}_K^{\mathrm{ab}} \longrightarrow \mathbb{F}_{p^2}^{\times},$$

where $\mathrm{G}_K^{\mathrm{ab}}$ is the Galois group of the maximal abelian extension K^{ab}/K.

For a prime $\mathfrak{L}$ of K, let $\mathcal{O}_{K,\mathfrak{L}}$ be the completion of $\mathcal{O}_K$ at $\mathfrak{L}$, and

$$r_p(\mathfrak{L}) : \mathcal{O}_{K,\mathfrak{L}}^{\times} \longrightarrow \mathbb{F}_{p^2}^{\times}$$

the composition

$$\mathcal{O}_{K,\mathfrak{L}}^{\times} \xrightarrow{\omega_{\mathfrak{L}}} \mathrm{G}_K^{\mathrm{ab}} \xrightarrow{\varrho_p^{\mathrm{ab}}} \mathbb{F}_{p^2}^{\times}.$$

Here $\omega_{\mathfrak{L}}$ is the Artin map.

Proposition 2.1 ([3, Proposition 4.7 (2)])**.** *If $\mathfrak{L} \nmid p$, then $r_p(\mathfrak{L})^{12} = 1$.*

Fix a prime $\mathfrak{P}$ of K above p. Then we have an isomorphism $\kappa(\mathfrak{P}) \cong \mathbb{F}_{p^{f_{\mathfrak{P}}}}$ of finite fields. Let $t_{\mathfrak{P}} := \gcd(2, f_{\mathfrak{P}}) \in \{1, 2\}$.

Proposition 2.2 ([3, Proposition 4.8])**.** (1) *There is a unique element $c_{\mathfrak{P}} \in \mathbb{Z}/(p^{t_{\mathfrak{P}}} - 1)\mathbb{Z}$ satisfying $r_p(\mathfrak{P})(u) = \mathrm{Norm}_{\kappa(\mathfrak{P})/\mathbb{F}_{p^{t_{\mathfrak{P}}}}}(\widetilde{u})^{-c_{\mathfrak{P}}}$ for any $u \in \mathcal{O}_{K,\mathfrak{P}}^{\times}$. Here, $\widetilde{u} \in \kappa(\mathfrak{P})$ is the reduction of u modulo $\mathfrak{P}$.*

(2) $\dfrac{2c_{\mathfrak{P}}}{t_{\mathfrak{P}}} \equiv e_{\mathfrak{P}} \bmod (p-1)$.

Corollary 2.3. *For any prime number $l \neq p$, we have $r_p(\mathfrak{P})(l^{-1})^2 = l^{e_{\mathfrak{P}} f_{\mathfrak{P}}} \bmod p$.*

Proof. $r_p(\mathfrak{P})(l^{-1})^2 = (\mathrm{Norm}_{\kappa(\mathfrak{P})/\mathbb{F}_{p^{t_{\mathfrak{P}}}}}(l^{-1})^{-c_{\mathfrak{P}}})^2 = \mathrm{Norm}_{\mathbb{F}_{p^{f_{\mathfrak{P}}}}/\mathbb{F}_{p^{t_{\mathfrak{P}}}}}(l)^{2c_{\mathfrak{P}}} = l^{\frac{2c_{\mathfrak{P}} f_{\mathfrak{P}}}{t_{\mathfrak{P}}}} = l^{e_{\mathfrak{P}} f_{\mathfrak{P}}} \bmod p$.

□

For a prime number l, the action of G_K on the l-adic Tate module $T_l A$ yields a representation

$$R_l : \mathrm{G}_K \longrightarrow \mathrm{Aut}_{\mathcal{O}}(T_l A) \cong \mathcal{O}_l^{\times} \subseteq B_l^{\times},$$

where $\mathrm{Aut}_{\mathcal{O}}(T_l A)$ is the group of automorphisms of $T_l A$ commuting with the action of $\mathcal{O}$, and $\mathcal{O}_l = \mathcal{O} \otimes_{\mathbb{Z}} \mathbb{Z}_l$, $B_l = B \otimes_{\mathbb{Q}} \mathbb{Q}_l$. Let $\mathrm{Nrd}_{B_l/\mathbb{Q}_l}$ be the reduced norm on B_l. Let $\mathfrak{M}$ be a prime of K, and $F_{\mathfrak{M}} \in \mathrm{G}_K$ a Frobenius element at $\mathfrak{M}$. For each $e \geq 1$, there is an integer $a(F_{\mathfrak{M}}^e) \in \mathbb{Z}$ satisfying

$$\mathrm{Nrd}_{B_l/\mathbb{Q}_l}(T - R_l(F_{\mathfrak{M}}^e)) = T^2 - a(F_{\mathfrak{M}}^e)T + (\mathrm{N}_{\mathfrak{M}})^e \in \mathbb{Z}[T]$$

for any l prime to $\mathfrak{M}$.

Proposition 2.4 ([3, Proposition 5.3]). (1) *We have* $a(F_{\mathfrak{M}}^e)^2 \leq 4(\mathrm{N}\mathfrak{M})^e$ *for any positive integer* e.

(2) *Assume* $\mathfrak{M} \nmid p$. *Then*

$$a(F_{\mathfrak{M}}^e) \equiv \varrho_p(F_{\mathfrak{M}}^e) + (\mathrm{N}\mathfrak{M})^e \varrho_p(F_{\mathfrak{M}}^e)^{-1} \bmod p$$

for any positive integer e.

Let $\alpha_{\mathfrak{M}}, \overline{\alpha}_{\mathfrak{M}} \in \mathbb{C}$ be the roots of $T^2 - a(F_{\mathfrak{M}})T + \mathrm{N}\mathfrak{M}$. Then $\alpha_{\mathfrak{M}} + \overline{\alpha}_{\mathfrak{M}} = a(F_{\mathfrak{M}})$ and $\alpha_{\mathfrak{M}}\overline{\alpha}_{\mathfrak{M}} = \mathrm{N}\mathfrak{M}$. We see that the roots of $T^2 - a(F_{\mathfrak{M}}^e)T + (\mathrm{N}\mathfrak{M})^e$ are $\alpha_{\mathfrak{M}}^e, \overline{\alpha}_{\mathfrak{M}}^e$. Then $\alpha_{\mathfrak{M}}^e + \overline{\alpha}_{\mathfrak{M}}^e = a(F_{\mathfrak{M}}^e)$. We have the following corollary to Proposition 2.4(1) (for $e = 1$):

Corollary 2.5. *We have* $a(F_{\mathfrak{M}}^e) \in \mathcal{C}(\mathrm{N}\mathfrak{M}, e)$ *for any positive integer* e.

For a later use, we give the following lemma:

Lemma 2.6. *Let* m *be the residue characteristic of* $\mathfrak{M}$. *The the following conditions are equivalent:*

(i) $m \mid a(F_{\mathfrak{M}})$.
(ii) $m \mid a(F_{\mathfrak{M}}^e)$ *for a positive integer* e.
(iii) $m \mid a(F_{\mathfrak{M}}^e)$ *for any positive integer* e.

Proof. For each $e \geq 1$, there is a polynomial $P_e(S,T) \in \mathbb{Z}[S,T]$ such that $(S+T)^e = S^e + T^e + STP_e(S+T, ST)$. Then $a(F_{\mathfrak{M}})^e = a(F_{\mathfrak{M}}^e) + \mathrm{N}\mathfrak{M}P_e(a(F_{\mathfrak{M}}), \mathrm{N}\mathfrak{M})$. Since $m \mid \mathrm{N}\mathfrak{M}$, we have $m \mid a(F_{\mathfrak{M}})$ if and only if $m \mid a(F_{\mathfrak{M}}^e)$.

□

3 Proof of the main result

Now we prove Theorem 1.1. Suppose that the assumption of Theorem 1.1 holds. Assume that there is a point $x \in M^B(k)$. When $B \otimes_{\mathbb{Q}} k \ncong \mathrm{M}_2(k)$, let K_0 be a quadratic extension of k satisfying $B \otimes_{\mathbb{Q}} K_0 \cong \mathrm{M}_2(K_0)$. Let

$$K := \begin{cases} k & \text{if } B \otimes_{\mathbb{Q}} k \cong \mathrm{M}_2(k), \\ K_0 & \text{if } B \otimes_{\mathbb{Q}} k \ncong \mathrm{M}_2(k). \end{cases}$$

Note that the degree $[K : \mathbb{Q}]$ is even. Then x is represented by a QM-abelian surface (A, i) by $\mathcal{O}$ over K (see [3, Theorem 1.1]). Since $B \notin \mathcal{S}(k, \mathfrak{q})$, there is a prime divisor p of $d(B)$ such that $p \neq q$ and p does not belong to

$$\begin{cases} \mathcal{P}(\mathcal{D}(\mathrm{N}\mathfrak{q}, e_{\mathfrak{q}})) & \text{if } B \otimes_{\mathbb{Q}} k \cong \mathrm{M}_2(k), \\ \mathcal{P}(\mathcal{D}(\mathrm{N}\mathfrak{q}, 2e_{\mathfrak{q}})) & \text{if } B \otimes_{\mathbb{Q}} k \ncong \mathrm{M}_2(k). \end{cases}$$

Fix such p, and let

$$\varrho_p : \mathrm{G}_K \longrightarrow \mathbb{F}_{p^2}^{\times}$$

be a canonical isogeny character at p associated to (A, i).

By Proposition 2.1, the character ϱ_p^{12} is unramified outside p. Then it is identified with a character $\mathfrak{I}_K(p) \longrightarrow \mathbb{F}_{p^2}^\times$, where $\mathfrak{I}_K(p)$ is the group of fractional ideals of K prime to p. When $B \otimes_{\mathbb{Q}} k \ncong \mathrm{M}_2(k)$, we may assume that $\mathfrak{q}$ is ramified in K/k by replacing K_0 if necessary. In any case, let $\mathfrak{Q}$ be the unique prime of K above $\mathfrak{q}$. Note that $\mathfrak{Q}$ is the unique prime of K above q, and so $q\mathcal{O}_K = \mathfrak{Q}^{e_\mathfrak{Q}}$, $(\mathrm{N}_\mathfrak{Q})^{e_\mathfrak{Q}} = (q^{f_\mathfrak{Q}})^{e_\mathfrak{Q}} = q^{[K:\mathbb{Q}]}$. Then by Corollary 2.3, we have

$$\varrho_p^{12}(F_\mathfrak{Q}^{e_\mathfrak{Q}}) = \varrho_p^{12}(\mathfrak{Q}^{e_\mathfrak{Q}}) = \varrho_p^{12}(q\mathcal{O}_K) = \varrho_p^{12}(1, \cdots, 1, q, \cdots, q, \cdots)$$

$$= \varrho_p^{12}(q^{-1}, \cdots, q^{-1}, 1, \cdots, 1, \cdots) = \prod_{\mathfrak{P}|p} r_p(\mathfrak{P})^{12}(q^{-1}) \equiv \prod_{\mathfrak{P}|p} q^{6e_\mathfrak{P} f_\mathfrak{P}} = q^{6[K:\mathbb{Q}]} \bmod p.$$

Here, $(1, \cdots, 1, q, \cdots, q, \cdots)$ (resp. $(q^{-1}, \cdots, q^{-1}, 1, \cdots, 1, \cdots)$) is the idèle of K whose components above p are 1 and the others q (resp. whose components above p are q^{-1} and the others 1), and $\mathfrak{P}$ runs through the primes of K above p. On the other hand, we have

$$a(F_\mathfrak{Q}^{e_\mathfrak{Q}}) \equiv \varrho_p(F_\mathfrak{Q}^{e_\mathfrak{Q}}) + (\mathrm{N}_\mathfrak{Q})^{e_\mathfrak{Q}} \varrho_p(F_\mathfrak{Q}^{e_\mathfrak{Q}})^{-1} = \varrho_p(F_\mathfrak{Q}^{e_\mathfrak{Q}}) + q^{[K:\mathbb{Q}]} \varrho_p(F_\mathfrak{Q}^{e_\mathfrak{Q}})^{-1} \bmod p$$

by Proposition 2.4(2). Let $\varepsilon := q^{-\frac{[K:\mathbb{Q}]}{2}} \varrho_p(F_\mathfrak{Q}^{e_\mathfrak{Q}}) \in \mathbb{F}_{p^2}^\times$. Then

$$\varepsilon^{12} = 1 \qquad \text{and} \qquad a(F_\mathfrak{Q}^{e_\mathfrak{Q}}) \equiv (\varepsilon + \varepsilon^{-1}) q^{\frac{[K:\mathbb{Q}]}{2}} \bmod p.$$

Therefore

$$a(F_\mathfrak{Q}^{e_\mathfrak{Q}}) \equiv 0, \pm q^{\frac{[K:\mathbb{Q}]}{2}}, \pm 2q^{\frac{[K:\mathbb{Q}]}{2}} \bmod p \qquad \text{or} \qquad a(F_\mathfrak{Q}^{e_\mathfrak{Q}})^2 \equiv 3q^{[K:\mathbb{Q}]} \bmod p.$$

By Corollary 2.5, we have $a(F_\mathfrak{Q}^{e_\mathfrak{Q}}) \in \mathcal{C}(\mathrm{N}_\mathfrak{Q}, e_\mathfrak{Q})$. We also have

$$\mathrm{N}_\mathfrak{Q} = \mathrm{N}_\mathfrak{q} \qquad \text{and} \qquad e_\mathfrak{Q} = \begin{cases} e_\mathfrak{q} & \text{if } B \otimes_{\mathbb{Q}} k \cong \mathrm{M}_2(k), \\ 2e_\mathfrak{q} & \text{if } B \otimes_{\mathbb{Q}} k \ncong \mathrm{M}_2(k). \end{cases}$$

Then

$$a(F_\mathfrak{Q}^{e_\mathfrak{Q}}), a(F_\mathfrak{Q}^{e_\mathfrak{Q}}) \pm q^{\frac{[K:\mathbb{Q}]}{2}}, a(F_\mathfrak{Q}^{e_\mathfrak{Q}}) \pm 2q^{\frac{[K:\mathbb{Q}]}{2}}, a(F_\mathfrak{Q}^{e_\mathfrak{Q}})^2 - 3q^{[K:\mathbb{Q}]} \in \mathcal{D}(\mathrm{N}_\mathfrak{Q}, e_\mathfrak{Q}).$$

Since $p \notin \mathcal{P}(\mathcal{D}(\mathrm{N}_\mathfrak{q}, e_\mathfrak{Q}))$, we have

(1) $a(F_\mathfrak{Q}^{e_\mathfrak{Q}}) = 0, \pm q^{\frac{[K:\mathbb{Q}]}{2}}, \pm 2q^{\frac{[K:\mathbb{Q}]}{2}}$, or

(2) $a(F_\mathfrak{Q}^{e_\mathfrak{Q}})^2 = 3q^{[K:\mathbb{Q}]}$.

[Case (1)]. In this case, we have $q \mid a(F_\mathfrak{Q}^{e_\mathfrak{Q}})$. Then by Lemma 2.6, we have $q \mid a(F_\mathfrak{Q})$. Since $f_\mathfrak{Q} (= f_\mathfrak{q})$ is odd, we obtain $B \otimes_{\mathbb{Q}} \mathbb{Q}(\sqrt{-q}) \cong \mathrm{M}_2(\mathbb{Q}(\sqrt{-q}))$ or ($q = 2$ and $B \otimes_{\mathbb{Q}} \mathbb{Q}(\sqrt{-1}) \cong \mathrm{M}_2(\mathbb{Q}(\sqrt{-1}))$) (see [3, Theorem 2.1, Propositions 2.3 and 5.1 (1)]). This contradicts $B \in \mathcal{B}(q)$.
[Case (2)]. In this case, $q = 3$ and $[K : \mathbb{Q}]$ is odd, which is a contradiction.

Therefore we conclude $M^B(k) = \emptyset$.

□

Table 1:

(N, e)	$\mathcal{C}(N, e)$	$\mathcal{D}(N, e)$	$\mathcal{P}(\mathcal{D}(N, e))$
$(2, 2)$	0, −3, −4	0, ±1, ±2, −3, ±4, −5, −6, −7, −8, −12	2, 3, 5, 7
$(2, 4)$	1, ±8	0, 1, −3, ±4, 5, −7, ±8, 9, ±12, ±16, −47	2, 3, 5, 7, 47
$(2, 6)$	0, 9, −16	0, 1, −7, ±8, 9, ±16, 17, −24, 25, −32, 64, −111, −192	2, 3, 5, 7, 17, 37
$(2, 8)$	−31, 32	0, 1, −15, 16, −31, 32, −47, 48, −63, 64, 193, 256	2, 3 , 5, 7, 31, 47, 193
$(2, 10)$	0, 57, −64	0, −7, 25, ±32, 57, ±64, 89, −96, 121, −128, 177, 1024, −3072	2, 3, 5, 7, 11, 19, 59, 89
$(2, 12)$	−47, ±128	0, 17, −47, ±64, 81, −111, ±128, −175, ±192, ±256, 4096, −10079	2, 3, 5, 7, 17, 37, 47, 10079
$(2, 14)$	0, −87, −256	0, 41, −87, ±128, 169, −215, ±256, −343, −384, −512, 16384,−41583, −49152	2, 3, 5, 7, 13, 29, 41, 43, 83, 167
$(2, 16)$	449, 512	0, −63, 193, 256, 449, 512, 705, 768, 961, 1024, 4993, 65536	2, 3, 5, 7, 31, 47, 193, 449, 4993
$(3, 2)$	−2, 3, −5, −6	0, 1, −2, ±3, 4, −5, ±6, −8, ±9, −11, −12, −18, −23	2, 3, 5, 11, 23
$(3, 4)$	7, −9, −14, 18	0, −2, 4, −5, 7, ±9, −11, −14, 16, ±18, −23, 25, ±27, −32, 36, −47, 81, −162, −194	2, 3, 5, 7, 11, 23, 47, 97
$(3, 6)$	10, 46, −54	0, −8, 10, −17, 19, −27, 37, −44, 46, −54, 64, −71, 73, −81, 100, −108, 729, −2087	2, 3, 5, 11, 17, 19, 23, 37, 71, 73, 2087
$(3, 8)$	34, −81, −113, 162	0, −32, 34, −47, 49, ±81, −113, 115, −128, ±162, −194, 196, ±243, −275, 324, 6561, −6914, −13122, −18527	2, 3, 5, 7, 11, 17, 23, 47, 97, 113, 191, 3457
$(3, 10)$	243, 475, −482, −486	0, 4, −11, 232, −239, ±243, 475, −482, ±486, 718, −725, ±729, 961, −968, −972, 48478, 55177, 59049, −118098	2, 3, 5, 11, 19, 23, 29, 31, 239, 241, 359, 2399, 24239
$(3, 12)$	658, −1358, 1458	0, −71, 100, −629, 658, 729, −800, −1358, 1387, 1458, −2087, 2116, 2187, −2816, 2916, 249841, 531441, −1161359	2, 3, 5, 7, 11, 17, 19, 23, 37, 47, 71, 73, 97, 433, 577, 1009, 1151, 2087
$(3, 14)$	2187, 2515, 3022, −4374	0, 328, 835, −1352, −1859, ±2187, 2515, 3022, ±4374, 4702, 5209, ±6561, 6889, 7396, −8748, 4782969, −5216423, −8023682, −9565938	2, 3, 5, 11, 13, 23, 41, 43, 83, 167, 337, 503, 673, 1511, 2351, 5209, 24023
$(3, 16)$	−353, −6561, −11966, 13122	0, −353, 1156, −5405, 6208, ±6561, −6914, −11966, 12769, ±13122, −13475, −18527, ±19683, −25088, 26244, 14044993, 43046721, −86093442, −129015554	2, 3, 5, 7, 11, 17, 23, 31, 47, 97, 113, 191, 193, 353, 383, 2113, 3457, 30529, 36671

4 Counterexamples to the Hasse principle

We have computed the sets $\mathcal{C}(N,e), \mathcal{D}(N,e), \mathcal{P}(\mathcal{D}(N,e))$ in several cases as seen in Table 1. Then we obtain the following counterexamples to the Hasse principle on M^B over number fields:

Proposition 4.1. (1) *Let $d(B) = 39$, and let $k = \mathbb{Q}(\sqrt{2}, \sqrt{-13})$ or $\mathbb{Q}(\sqrt{-2}, \sqrt{-13})$. Then $B \otimes_{\mathbb{Q}} k \cong \mathrm{M}_2(k)$, $M^B(k) = \emptyset$ and $M^B(k_v) \neq \emptyset$ for any place v of k. Here, k_v is the completion of k at v.*

(2) *Let L be the subfield of $\mathbb{Q}(\zeta_9)$ satisfying $[L : \mathbb{Q}] = 3$, and let $(d(B), k) = (62, L(\sqrt{-39}))$ or $(86, L(\sqrt{-15}))$. Then $B \otimes_{\mathbb{Q}} k \ncong \mathrm{M}_2(k)$, $M^B(k) = \emptyset$ and $M^B(k_v) \neq \emptyset$ for any place v of k.*

Proof. (1) The prime number 3 (resp. 13) is inert (resp. ramified) in $\mathbb{Q}(\sqrt{-13})$. Then $B \otimes_{\mathbb{Q}} \mathbb{Q}(\sqrt{-13}) \cong \mathrm{M}_2(\mathbb{Q}(\sqrt{-13}))$, and so $B \otimes_{\mathbb{Q}} k \cong \mathrm{M}_2(k)$.

Applying Theorem 1.1 to $q = 2$, we obtain $M^B(k) = \emptyset$. In fact, $(e_{\mathfrak{q}}, f_{\mathfrak{q}}) = (4, 1)$ where $\mathfrak{q}$ is the unique prime of k above $q = 2$, and the prime divisor 13 of $d(B)$ does not belong to $\mathcal{P}(\mathcal{D}(2,4)) \cup \{2\}$ (see Table 1). Since 3 (resp. 13) splits in $\mathbb{Q}(\sqrt{-2})$ (resp. $\mathbb{Q}(\sqrt{-1})$), we have $B \otimes_{\mathbb{Q}} \mathbb{Q}(\sqrt{-2}) \ncong \mathrm{M}_2(\mathbb{Q}(\sqrt{-2}))$ (resp. $B \otimes_{\mathbb{Q}} \mathbb{Q}(\sqrt{-1}) \ncong \mathrm{M}_2(\mathbb{Q}(\sqrt{-1}))$).

By [3, p.94], we have $M^B(\mathbb{Q}(\sqrt{-13})_w) \neq \emptyset$ for any place w of $\mathbb{Q}(\sqrt{-13})$ (cf. [4]). Therefore $M^B(k_v) \neq \emptyset$ for any place v of k.

(2) For a field F of characteristic $\neq 2$ and two elements $a, b \in F^\times$, let

$$\left(\frac{a,b}{F}\right) = F + Fe + Ff + Fef$$

be the quaternion algebra over F defined by

$$e^2 = a,\ f^2 = b,\ ef = -fe.$$

For a prime number p, let e_p, f_p, g_p be the ramification index of p in $k/\mathbb{Q}$, the degree of the residue field extension above p in $k/\mathbb{Q}$, and the number of primes of k above p respectively.

Let $(d(B), k) = (62, L(\sqrt{-39}))$ (resp. $(86, L(\sqrt{-15}))$). First, we prove $B \otimes_{\mathbb{Q}} k \ncong \mathrm{M}_2(k)$. We see $B \cong \left(\frac{62, 13}{\mathbb{Q}}\right)$ (resp. $\left(\frac{86, 5}{\mathbb{Q}}\right)$) by [6, §3.6 g)]. We have $(e_2, f_2, g_2) = (1, 3, 2)$. Let v be place of k above 2. By the same argument as in the proof of [1, Proposition 8.1], we have $B \otimes_{\mathbb{Q}} k_v \ncong \mathrm{M}_2(k_v)$. Therefore $B \otimes_{\mathbb{Q}} k \ncong \mathrm{M}_2(k)$.

Applying Theorem 1.1 to $q = 3$, we obtain $M^B(k) = \emptyset$. In fact, $(e_{\mathfrak{q}}, f_{\mathfrak{q}}) = (6, 1)$ where $\mathfrak{q}$ is the unique prime of k above $q = 3$, and the prime divisor 31 (resp. 43) of $d(B)$ does not belong to $\mathcal{P}(\mathcal{D}(3, 12)) \cup \{3\}$. Since 31 (resp. 43) splits in $\mathbb{Q}(\sqrt{-3})$, we have $B \otimes_{\mathbb{Q}} \mathbb{Q}(\sqrt{-3}) \ncong \mathrm{M}_2(\mathbb{Q}(\sqrt{-3}))$.

By [5, Table 1], we have $M^B(\mathbb{Q}(\sqrt{-39})_w) \neq \emptyset$ (resp. $M^B(\mathbb{Q}(\sqrt{-15})_w) \neq \emptyset$) for any place w of $\mathbb{Q}(\sqrt{-39})$ (resp. $\mathbb{Q}(\sqrt{-15})$). Therefore $M^B(k_v) \neq \emptyset$ for any place v of k.

□

References

[1] *K. Arai*, Algebraic points on Shimura curves of $\Gamma_0(p)$-type (III), preprint, available at the web page (http://arxiv.org/pdf/1303.5270.pdf).

[2] *K. Buzzard*, Integral models of certain Shimura curves, Duke Math. J. **87** (1997), no. 3, 591–612.

[3] *B. Jordan*, Points on Shimura curves rational over number fields, J. Reine Angew. Math. **371** (1986), 92–114.

[4] *B. Jordan, R. Livné*, Local Diophantine properties of Shimura curves, Math. Ann. **270** (1985), no. 2, 235–248.

[5] *V. Rotger, C. de Vera-Piquero*, Galois representations over fields of moduli and rational points on Shimura curves, Canad. J. Math. **66** (2014), 1167–1200.

[6] *H. Shimizu*, Hokei kansū. I–III (Japanese) [Automorphic functions. I–III] Second edition. Iwanami Shoten Kiso Sūgaku [Iwanami Lectures on Fundamental Mathematics], 8. Daisū [Algebra], vii. Iwanami Shoten, Tokyo, 1984.

[7] *G. Shimura*, On the real points of an arithmetic quotient of a bounded symmetric domain, Math. Ann. **215** (1975), 135–164.

(Keisuke Arai) Department of Mathematics, School of Engineering, Tokyo Denki University, 5 Senju Asahi-cho, Adachi-ku, Tokyo 120-8551, Japan

E-mail address: araik@mail.dendai.ac.jp